

A C^0 -ESTIMATE FOR THE PARABOLIC MONGE-AMPÈRE EQUATION ON COMPLETE NON-COMPACT KÄHLER MANIFOLDS

ALBERT CHAU¹ AND LUEN-FAI TAM²

ABSTRACT. In this article we study the Kähler Ricci flow, the corresponding parabolic Monge Ampère equation and complete non-compact Kähler Ricci flat manifolds. In our main result Theorem 1 we prove that if (M, g) is sufficiently close to being Kähler Ricci flat in a suitable sense, then the Kähler-Ricci flow (1) has a long time smooth solution $g(t)$ converging smoothly uniformly on compact sets to a complete Kähler Ricci flat metric on M . The main step is to obtain a uniform C^0 -estimates for the corresponding parabolic Monge Ampère equation. Our results on this can be viewed as a parabolic version of the main results in [10] on the elliptic Monge Ampère equation.

1. Let (M^n, g_0) be a complete non-compact Kähler manifold with complex dimension n and consider the following Kähler-Ricci flow on M :

$$(1) \quad \begin{cases} \frac{\partial g_{i\bar{j}}}{\partial t} = -R_{i\bar{j}} \\ g_{i\bar{j}}(x, 0) = (g_0)_{i\bar{j}} \end{cases}$$

We are interested in studying when (1) admits a long time solution $g(t)$ converging smoothly on M to a complete Kähler metric $g(\infty)$. Such a limit $g(\infty)$ must be Kähler Einstein with zero scalar curvature by (1). We are thus interested in studying when (M^n, g_0) converges to a Kähler Ricci flat metric under (1). When M is compact Cao [3] established that a necessary and sufficient condition for such convergence is that:

$$(2) \quad (R_0)_{i\bar{j}} = (f_0)_{i\bar{j}}$$

Date: July 2008.

¹Research partially supported by NSERC grant no. #327637-06.

²Research partially supported by Hong Kong RGC General Research Fund #GRF 2160357.

for a smooth potential function f_0 on M , thus re-establishing the famous Calabi Conjecture first proved by Yau [11]. In Theorem 1 we establish a non-compact version of Cao's result. We prove that when (M^n, g_0) is complete, non-compact with bounded curvature, with volume growth $V_{x_0}(r) \leq Cr^{2n}$ for some x_0 and C for all r , and satisfies a Sobolev inequality, then:

Under the above conditions, the Kähler-Ricci flow (1) has a long time solution $g(t)$ converging smoothly on M provided (2) is satisfied and $|f_0|(x) \leq \frac{C}{1+\rho_0^{2+\epsilon}(x)}$ for some $C, \epsilon > 0$ and all x .

See Theorem 1 for details. The result is motivated by the work of Tian-Yau [10] who proved the existence of a Kähler Ricci flat metric in the complement of a divisor in a projective variety under certain conditions. They first constructed a Kähler metric satisfying the condition in Theorem 1 and then solve the elliptic complex Monge Ampère equation to obtain a Kähler Ricci flat metric. Our results here can be viewed as a parabolic version of the results on the elliptic Monge Ampère equation in [10]. We also refer to [1, 3] for results on convergence of the Kähler-Ricci flow to Kähler Einstein metrics with negative scalar curvature. In [3], it was proved that (1) converges after rescaling to a Kähler Einstein metric with negative scalar curvature provided $(R_0)_{i\bar{j}} + (g_0)_{i\bar{j}} = (f_0)_{i\bar{j}}$ for smooth f_0 . A non-compact version of this result was proved in [1].

2. Let (M^n, g_0) be a complete non-compact Kähler manifold with complex dimension n such that (2) holds for some smooth potential f_0 on M . We want to study the long time behavior of the Kähler-Ricci flow (1), when f_0 is sufficiently close to zero in a suitable sense. Note that (M^n, g_0) is Kähler Einstein with zero scalar curvature when $f_0 = 0$. We are thus interested in the behavior of the Kähler-Ricci flow on complete Kähler manifolds which are close to being Kähler Einstein. We will prove the following:

Theorem 1. *Let (M^n, g_0) be a complete non-compact Kähler manifold with bounded curvature and $n \geq 3$. Assume the following:*

(a) *Condition (2) holds for smooth f_0 satisfying:*

$$(3) \quad |f_0|(x) \leq \frac{C_1}{1 + \rho_0^{2+\epsilon}(x)}$$

for some $C_1, \epsilon > 0$, and all $x \in M$ where $\rho_0(x)$ is the distance function from a fixed $p \in M$

(b) *The following Sobolev inequality is true:*

$$(4) \quad \left(\int_M |\phi|^{\frac{2n}{n-1}} dV_0 \right)^{\frac{n-1}{n}} \leq C_2 \int_M |\nabla_0 \phi|^2$$

for some $C_2 > 0$ and all $\phi \in C_0^\infty(M)$.

(c) *There exists constant $C_3 > 0$ such that*

$$(5) \quad V_0(r) \leq C_3 r^{2n}$$

for some $C_3 > 0$ and all r where $V_0(r)$ is the volume of the geodesic ball with radius r centered at some $p \in M$.

Then (1) has long time solution $g(t)$ converging uniformly on compact sets in the C^∞ topology to a complete Kähler Ricci flat metric g_∞ on M .

We will also consider the following parabolic Monge Ampère equation corresponding to (1):

$$(6) \quad \begin{cases} \frac{\partial u}{\partial t} = \log \frac{\det((g_0)_{k\bar{l}} + u_{k\bar{l}})}{\det((g_0)_{k\bar{l}})} - f_0 \\ u(x, 0) = 0 \end{cases}$$

The relationship between the two equations can be described simply as follows. If (6) has a smooth solution u on $M \times [0, T)$ then $g_{i\bar{j}} = (g_0)_{i\bar{j}} + u_{i\bar{j}}$ is a smooth solution to (1). Conversely, if (1) has a solution g then

$$u(x, t) = \int_0^t \log \frac{\det(g_{i\bar{j}})}{\det((g_0)_{i\bar{j}})}(x, s) ds - t f_0(x)$$

is a solution to (6) as shown in [1].

Lemma 1. *Let (M^n, g_0) be a complete non-compact Kähler manifold with bounded curvature such that (2) holds for a smooth bounded potential f_0 . Suppose $g(t)$ is a smooth solution to (1) on $M \times [0, T)$ such that $g(t)$ has bounded curvature for each $0 < t < T$. Then if $u(x, t)$ is the corresponding solution to (6), then $f(t) = -u_t$ satisfies the following for each $0 < t < T$:*

- (i) $|f|$, $t|\nabla_t f|^2$ and $t\Delta_t f$ are all uniformly bounded on M by a constant depending only on f_0 and g_0 .
- (ii) All the covariant derivatives of $f(t)$ relative to $g(t)$ are bounded on M .

- (iii) *If in addition, $|f_0(x)| \leq C/(1 + \rho_0(x))^N$ for some $N > 1$ and distance function ρ_0 relative some $p \in M$, then there is a constant C_t such that $|f(x, t)| \leq C_t/(1 + \rho_t(x))^N$ where $\rho_t(x)$ is the distance relative to p with respect to $g(t)$.*

Proof. Fix some $t_0 \in (0, T)$. We begin by noting that by (1) and the uniform curvature bounds, $g(t)$ is uniformly equivalent to g_0 on $M \times [0, t_0]$. Moreover, by [8] all covariant derivatives of $g(t)$ are bounded on $M \times (0, t_0]$. In particular, it follows that $(M, g(t))$ has bounded geometry of all orders for each $t \in (0, t_0]$.

Now differentiating (6) gives

$$(7) \quad \left(\frac{\partial}{\partial t} - \Delta_t \right) f = 0.$$

As f_0 is bounded, it follows from the above bounds on $g(t)$ (and curvature) and the maximum principle in [8, Lemma 4.7] that $\sup_M |f(t)| \leq \sup_M |f_0|$ for all $t \in [0, t_0]$. From this, the fact that $\Delta_t f(t) = R(t)$, the bounded geometry of $g(t)$ mentioned above and standard elliptic theory, we see that (ii) is true since $t_0 \in (0, T)$ was arbitrarily chosen. Part (i) now follows from the uniform bound on $|f(t)|$, the bounded geometry of $g(t)$ mentioned above, the main Theorem in [4] and by letting $t_0 \rightarrow T$.

We now prove (iii). By Lemma 4.5 in [8] there exists a smooth exhaustion function ρ_0 such that for some positive constants K_1, K_2, K_3 , depending only on n, t_0 and the uniform bound on $Rm(x, t)$ on $M \times [0, t_0]$, satisfying

- (1) $K_1(r_t + 1) \leq \rho_0 \leq K_2(r_t + 1)$
- (2) $|\nabla_t \rho_0|, |\nabla_t^2 \rho_0| \leq K_3$.

for all $t \in [0, t_0]$. Direct computation gives

$$\begin{aligned}
 (8) \quad & \frac{d}{dt}(\rho_0^{2N} f^2) \leq \rho_0^{2N} \Delta_t f^2 \\
 & = \Delta_t(\rho_0^{2N} f^2) - (\Delta_t \rho_0^{2N}) f^2 - 2\langle \nabla_t \rho_0^{2N}, \nabla_t f^2 \rangle_t \\
 & = \Delta_t(\rho_0^{2N} f^2) - 2N \rho_0^{2N-1} (\Delta_t \rho_0) f^2 \\
 & \quad - 2N(2N-1) \rho_0^{2N-2} |\nabla_t \rho_0|_t^2 f^2 - 4N \rho_0^{2N-1} \langle \nabla_t \rho_0, \nabla_t f^2 \rangle_t \\
 & \leq \Delta_t(\rho_0^{2N} f^2) + C_1 \rho_0^{2N} f^2 \\
 & \quad - 4N \rho_0^{2N-1} \langle \nabla_t \rho_0, \left(\frac{\nabla_t(\rho_0^{2N} f^2) - \nabla_t(\rho_0^{2N}) f^2}{\rho_0^{2N}} \right) \rangle_t \\
 & \leq \Delta_t(\rho_0^{2N} f^2) + C_2 \rho_0^{2N} f^2 - 4N \rho_0^{-1} \langle \nabla_t \rho_0, \nabla_t(\rho_0^{2N} f^2) \rangle_t
 \end{aligned}$$

for constants $C_1, C_2 > 0$ depending only on n, N, t_0 and the bound on $Rm(x, t)$ on $M \times [0, t_0]$.

On the other hand, for any $t \in [0, t_0]$ and for any $R > 0$, let φ be a cut off function on $B_0(R)$ which is zero outside $B_0(2R)$, and $|\nabla_0 \varphi|_0 \leq C/R$ for some constant C independent of t_0 and R . Then

$$(9) \quad \begin{aligned} \int_M \frac{\partial}{\partial t} (\varphi^2 f^2) dV_t &= \int_M \varphi^2 (\Delta_t f^2 dV_t - 2|\nabla_t f|_t^2) dV_t \\ &\leq 2 \int_M |\nabla_t \varphi|_t^2 f^2 dV_t + 2 \int_M \phi \Delta_t \phi f^2 dV_t \\ &\quad - \int_M \varphi^2 |\nabla_t f|_t^2 dV_t \end{aligned}$$

Integrating from 0 to t_0 and using the uniform bound on $\sup_M |f(t)|$, we have

$$(10) \quad \int_0^{t_0} \int_{B_0(R)} \varphi^2 |\nabla_t f|_t^2 dV_t dt \leq C_3 V_t(B_0(R)) \leq C_4 \exp(aR^2)$$

for some constants C_3, C_4 depending only on the bound on $Rm(x, t)$ on $M \times [0, t_0]$, n and t_0 . In particular, we can find $b > 0$ such that

$$(11) \quad \int_0^{t_0} \int_M \exp(-b\rho_t^2) |\nabla_t(\rho_0^{2N} f^2)|_t^2 dV_t dt < \infty.$$

Let $h = \exp(-C_2 t) \rho_0^{2N} f^2$, then by (8)

$$(12) \quad \frac{\partial}{\partial t} h \leq \Delta_t h - 4N \rho_0^{-1} \langle \nabla_t \rho_0, \nabla_t h \rangle_t.$$

Since at $t = 0$, h is bounded, if we let $h_1 = h - \sup_M h(0)$, then h_1 still satisfies (12) and $h_1(0) \leq 0$. By Theorem 4.3 in [5], together with (11), (12), and the properties of ρ_0 , we conclude that $h(t) \leq 0$ for all $t \in [0, t_0]$. Part (iii) of the lemma now follows by letting $t_0 \rightarrow T$. Note that we have used the fact that g and $g(t)$ are uniformly equivalent.

This completes the proof of the lemma. $\square$

Corollary 1. *Let (M^n, g_0) be a complete non-compact Kähler manifold with bounded curvature such that (2) is satisfied for a smooth bounded potential f_0 . Then (1) has a long time smooth solution.*

Proof. By [8], (1) has a solution on $M \times [0, T)$ where $T > 0$ is the maximal time satisfying the assumptions in Lemma 1. Let u and f as in the lemma. Then $|f(t)| \leq \sup_M |f_0|$ and hence $|u(x, t)| \leq t \sup_M |f_0|$ for all $(x, t) \in M \times [0, T)$ by Lemma 1. Hence $\sup_M u(x, t)$ cannot blowup in finite time. By the a priori estimates in [1], we conclude that $T = \infty$ and thus (1) has a long time smooth solution. $\square$

Note the solution $g(t)$ in the corollary satisfies the assumptions in Lemma 1 for any finite $t > 0$, and hence we may assume in Theorem 1 that f_0 has uniformly bounded covariant derivatives of all orders.

With the assumptions as in Theorem 1, let $g(t)$ be the long time solution as in Corollary 1 and let $u(x, t)$ and $f(x, t)$ be defined as before. In order to prove Theorem 1, we need the following lemmas.

Lemma 2. *Let (M, g_0) be a complete non-compact Kähler manifold with bounded curvature satisfying conditions (a) and (c) in Theorem 1 with $n \geq 3$ and let $g(t)$ be the longtime solution of the Kähler Ricci flow (1). For any $p_0 > n$ with $(p_0 + 1)(2 + \epsilon)/(n + p_0) > 2$ and $p^* = n(p_0 + 1)/(n + p_0) > 2$, there is a positive constant C independent of t such that*

$$\int_M |f(t)|^{p^*} dV_t \leq C$$

for all t .

Lemma 3. *Let (M, g_0) be as in Theorem 1 and let $g(t)$ be the longtime solution of the Kähler-Ricci flow (1). Then there is a positive constant C such that and*

$$\sup_{M \times [0, \infty)} |u| \leq C.$$

for all t .

Assuming the lemmas are true. Then we can prove Theorem 1 as follows:

Proof of Theorem 1. By Lemmas 2 and 3 and the a priori estimates in [1], we conclude that for all $t \geq 0$, $g(t)$ is uniformly equivalent to g_0 independent of t . Moreover, for any sequence $t_k \rightarrow \infty$ some subsequence of $u(x, t_k)$ (which we still denote by $u(x, t_k)$) converges in the C^∞ sense on compact subsets of M to a smooth limit v on M . Thus by Lemma 1 (i), we conclude that $\frac{\partial u}{\partial t}(t_k) = -f(t_k)$ converges uniformly on M to a constant c . Lemma 2 and the fact that M has infinite volume implies that c must be zero. Hence $g_{i\bar{j}} + v_{i\bar{j}}$ is a smooth complete Ricci flat Kähler metric on M . The proof of the theorem is complete once we show that the limit metric $g_{i\bar{j}} + v_{i\bar{j}}$ is independent of the t_k . This last fact follows by the next lemma. $\square$

The following result was basically proved in [2].

Lemma 4. *Let g and h be two equivalent complete Kähler metrics on a non-compact complex manifold M such that: (i) $h_{i\bar{j}} = g_{i\bar{j}} + v_{i\bar{j}}$ for some*

smooth bounded function v ; (ii) g has nonnegative Ricci curvature; and (iii)

$$\log \frac{\det(g_{i\bar{j}} + v_{i\bar{j}})}{\det(g_{i\bar{j}})} = 0.$$

Then $g = h$.

Proof. We sketch the proof. Since

$$\begin{aligned} 0 &= \int_0^1 \frac{\partial}{\partial s} \log \det(g_{i\bar{j}} + sv_{i\bar{j}}) \\ &= \left(\int_0^1 g^{i\bar{j}}(s) ds \right) v_{i\bar{j}} \end{aligned}$$

where $g_{i\bar{j}}(s) = g_{i\bar{j}} + sv_{i\bar{j}}$. Hence v satisfies $a^{i\bar{j}}v_{i\bar{j}} = 0$ for some Kähler metric $a_{i\bar{j}}$ which is uniformly equivalent to g . As g has nonnegative Ricci curvature, v must be constant by [6, 7]. $\square$

We now prove Lemma 2.

Lemma 2. Let $p_0 > n$ be such that

$$\frac{p_0 + 1}{n + p_0} (2 + \epsilon) > 2$$

and $p^* = \frac{n(p_0+1)}{n+p_0} > 2$.

By the assumption on f_0 , the volume growth of (M, g_0) and Lemma 1, we see that for any $T > 0$ and $p \geq p^*$, there is a constant C

$$(13) \quad \int_M |f(t)|^p dV_t \leq C$$

for all $0 \leq t \leq T$. Here we have used the fact that for any finite T , $g(t)$ is uniformly equivalent to g for $0 \leq t \leq T$.

For any even integer $p > p^* + 2$, we have

$$\begin{aligned} (14) \quad \frac{d}{dt} \int_M |f|^p dV_t &= \frac{d}{dt} \int_M f^p dV_t \\ &= p \int_M f^{p-1} \Delta_t f dV_t - \int_M f^p \Delta_t f dV_t \\ &= -p(p-1) \int_M f^{p-2} |\nabla_t f|_t^2 dV_t + p \int_M f^{p-1} |\nabla_t f|_t^2 dV_t \\ &\leq -(p(p-1) - pC_1) \int_M f^{p-2} |\nabla_t f|_t^2 dV_t. \end{aligned}$$

where $C_1 = \sup_M |f_0|$ and we have used the fact that for any t , $|\nabla f|_t$ is bounded on M and the fact that $p-2 > p^*$ to justify the integration by parts. Hence if $p \geq 1 + C_1$, then

$$(15) \quad \int_M |f(t)|^p dV_t \leq \int_M |f_0|^p dV_0 < \infty$$

We have to improve (15).

Let $v = \max\{f, 0\}$ and let φ be a nonnegative cutoff function on M independent of t . For any $p \geq p^* - 1 > 1$,

$$(16) \quad \begin{aligned} & \int_0^T \int_M \varphi^2 v^p \frac{\partial}{\partial t} f dV_t dt \\ &= \int_0^T \int_M \varphi^2 v^p \Delta_t f dV_t dt \\ &= - \int_0^T \int_M p v^{p-1} \varphi^2 |\nabla_t v|_t^2 dV_t dt - 2 \int_0^T \int_M v^p \varphi \langle \nabla_t \varphi, \nabla_t v \rangle_t dV_t dt \\ &\leq \frac{1}{p} \int_0^T \int_M v^{p+1} |\nabla_t \varphi|_t^2 dV_t dt. \end{aligned}$$

On the other hand,

$$(17) \quad \begin{aligned} & \int_0^T \int_M \varphi^2 v^p \frac{\partial f}{\partial t} dV_t dt \\ &= - \int_M \varphi^2 \int_0^T v^p \left(\frac{\partial}{\partial v t} \right) e^{f_0 - f} dt dV_0 \\ &= \frac{1}{p+1} \int_M \varphi^2 \int_0^T \frac{\partial}{\partial t} [v^{p+1} e^{f_0 - f}] dt dV_0 \\ &\quad + \frac{1}{p+1} \int_M \varphi^2 \int_0^T v^{p+1} e^{f_0 - f} \frac{\partial f}{\partial t} dt dV_0. \end{aligned}$$

Combining this with (16), we have

$$\begin{aligned}
 (18) \quad \int_M \varphi^2 v^{p+1} e^{f_0-f} dV_t|_{t=T} &\leq \int_M \varphi^2 v^{p+1} dV_t|_{t=0} + \frac{p+1}{p} \int_0^T \int_M v^{p+1} |\nabla_t \varphi|_t^2 dV_t dt \\
 &\quad - \int_M \varphi^2 e^{f_0} \int_0^T v^{p+1} \sum_{k=0}^{\infty} \frac{(-f)^k}{k!} \frac{\partial v}{\partial t} dt dV_0 \\
 &= \int_M \varphi^2 v^{p+1} dV_t|_{t=0} + \frac{p+1}{p} \int_0^T \int_M v^{p+1} |\nabla_t \varphi|_t^2 dV_t dt \\
 &\quad - \int_M \varphi^2 e^{f_0} \sum_{k=0}^{\infty} \frac{(-1)^k v^{p+k+2}}{k!(p+k+2)} dV_0|_{t=T} \\
 &\quad + \int_M \varphi^2 e^{f_0} \sum_{k=0}^{\infty} \frac{(-1)^k v^{p+k+2}}{k!(p+k+2)} dV_0|_{t=0}.
 \end{aligned}$$

Now

$$\begin{aligned}
 (19) \quad & - \int_M \varphi^2 e^{f_0} \sum_{k=0}^{\infty} \frac{(-1)^k v^{p+k+2}}{k!(p+k+2)} dV_0|_{t=T} + \int_M \varphi^2 e^{f_0} \sum_{k=0}^{\infty} \frac{(-1)^k v^{p+k+2}}{k!(p+k+2)} dV_0|_{t=0} \\
 &= - \int_M \varphi^2 v^{p+2} e^{f_0} \sum_{k=0}^{\infty} \frac{(-1)^k v^k}{k!(p+k+2)} dV_0|_{t=T} \\
 &\quad + \int_M \varphi^2 v^{p+2} e^{f_0} \sum_{k=0}^{\infty} \frac{(-1)^k v^k}{k!(p+k+2)} dV_0|_{t=0} \\
 &\leq C_2 \int_M v^{p+2} dV_t|_{t=T} + C_3 \int_M v^{p+2} dV_t|_{t=0} \\
 &\leq C_2 \int_M v^{p+2} dV_t|_{t=T} + C_4(p)
 \end{aligned}$$

where C_2, C_3, C_4 are independent of T . Here we have used the fact that $f(t)$ are uniformly bounded in spacetime. Combine (18) and (19) to give

$$\begin{aligned}
 (20) \quad \int_M \varphi^2 v^{p+1} dV_t|_{t=T} &\leq C_5 + \frac{p+1}{p} \int_0^T \int_M v^{p+1} |\nabla_t \varphi|_t^2 dV_t dt \\
 &\quad + C_2 \int_M v^{p+2} dV_t|_{t=T}
 \end{aligned}$$

for some constant independent of T . Letting φ approach the constant function 1 gives

$$(21) \quad \int_M v^{p+1} dV_t|_{t=T} \leq C_5 + C_2 \int_M v^{p+2} dV_t|_{t=T}.$$

Similarly one can prove that if $w = \max\{-f, 0\}$, then

$$(22) \quad \int_M w^{p+1} dV_t|_{t=T} \leq C_5 + C_2 \int_M w^{p+2} dV_t|_{t=T}$$

by modifying C_5 and C_2 if necessary, while still independent of T . Hence we have

$$(23) \quad \int_M |f|^{p+1} dV_t|_{t=T} \leq 2C_5 + C_2 \int_M |f|^{p+2}|_{t=T}$$

for all $p \geq p^* - 1$. By iteration and (15), we conclude that

$$(24) \quad \int_M |f|^{p^*} dV_t|_{t=T} \leq C_5$$

for some constant C_5 independent of T . $\square$

Proof of Lemma 3. We may now proceed as in [10] using (24). In particular note that for all t , $|u|$ also decays like $r_t^{-2-\epsilon}$ by Lemma 1 and hence we may integrate by parts as in [3], using (24) and the Sobolev inequality (4), to obtain the following for $p \geq p_0 > p^*$: (here p_0 and p^* are as in Lemma 2)

$$(25) \quad \begin{aligned} \left(\int_M |u|^{(p+1)\kappa} dV_0 \right)^{\frac{1}{\kappa}} &\leq C_6 p \int_M |u|^p |e^{f_0-f} - 1| dV_0 \\ &\leq C_7 p \int_M |u|^p |f_0 - f| dV_0 \\ &\leq C_7 p \left(\int_M |u|^{p+1} dV_0 \right)^{\frac{p}{p+1}} \left(\int_M |f_0 - f|^{p+1} dV_0 \right)^{\frac{1}{p+1}} \\ &\leq C_8 p \left(\int_M |u|^{p+1} dV_0 \right)^{\frac{p}{p+1}} \\ &\leq C_8 p \left(\int_M |u|^{p+1} dV_0 + 1 \right). \end{aligned}$$

Here C_8 is a constant independent of t, p and $\kappa = n/(n-1) > 1$ and we have used (24) and the fact that $f(t)$ is uniformly bounded on space

and time. Take $p = p_0$, we also have:

$$\begin{aligned}
 (26) \quad & \left(\int_M |u|^{(p_0+1)\kappa} dV_0 \right)^{\frac{1}{\kappa}} \\
 & \leq C_7 p_0 \int_M |u|^{p_0} |f_0 - f| dV_0 \\
 & \leq C_7 p_0 \left(\int_M |u|^{(p_0+1)\kappa} dV_0 \right)^{\frac{p_0}{(p_0+1)\kappa}} \left(\int_M |f_0 - f|^{\frac{(p_0+1)\kappa}{(p_0+1)\kappa - p_0}} dV_0 \right)^{1 - \frac{p_0}{(p_0+1)\kappa}} \\
 & = C_7 p_0 \left(\int_M |u|^{(p_0+1)\kappa} dV_0 \right)^{\frac{p_0}{(p_0+1)\kappa}} \left(\int_M |f_0 - f|^{p^*} dV_0 \right)^{\frac{1}{p^*}}.
 \end{aligned}$$

Hence by Lemma 2 we have

$$(27) \quad \left(\int_M |u|^{(p_0+1)\kappa} dV_0 \right)^{\frac{1}{(p_0+1)\kappa}} \leq C_9$$

for some constant C_9 independent of t . By (25) and Young's inequality we have that for $p \geq p_0$:

$$\begin{aligned}
 (28) \quad & \int_M |u|^{(p+1)\kappa} dV_0 + 1 \leq (C_8 p)^\kappa \left[\int_M |u|^{p+1} dV_0 + 1 \right]^\kappa + 1 \\
 & \leq (C_8 p)^\kappa \left[\int_M |u|^{p+1} dV_0 + 2 \right]^\kappa \\
 & \leq (2C_8 p)^\kappa \left[\int_M |u|^{p+1} dV_0 + 1 \right]^\kappa.
 \end{aligned}$$

Hence we have

$$\begin{aligned}
 (29) \quad & \left[\int_M |u|^{(p+1)\kappa} dV_0 + 1 \right]^{\frac{1}{\kappa(p+1)}} \leq (2C_8 p)^{\frac{1}{p+1}} \left[\int_M |u|^{p+1} dV_0 + 1 \right]^{\frac{1}{p+1}} \\
 & \leq (2C_8(p+1))^{\frac{1}{p+1}} \left[\int_M |u|^{p+1} dV_0 + 1 \right]^{\frac{1}{p+1}},
 \end{aligned}$$

That is to say, for all $q \geq p_0 + 1$,

$$(30) \quad \left[\int_M |u|^{\kappa q} dV_0 + 1 \right]^{\frac{1}{\kappa q}} \leq (2C_8 q)^{\frac{1}{q}} \left[\int_M |u|^q dV_0 + 1 \right]^{\frac{1}{q}}.$$

By iteration (see [10]), it is straight forward to show that

$$\sup_M |u| \leq C_{10} \left[\int_M |u|^{(p_0+1)\kappa} dV_0 + 1 \right]^{\frac{1}{(p_0+1)\kappa}} \leq C_{11}$$

for some constants $C_{10}, C_{11} > 0$ independent of t . Here we have used (27). This completes the proof of the lemma. $\square$

REFERENCES

1. Chau, A., *Convergence of the Kähler Ricci flow on non-compact Kähler Manifolds*, J. Differential Geom. 66 (2004), no. 2, 211–232.
2. Chau, A., *Stablility of the Kähler-Ricci flow at complete noncompact Kähler Einstein metrics*, Comtemporary Mathematics v. 367 (2005), 43–62.
3. Cao, Huai-Dong, *Deformation of Kähler metrics to Kahler Einstein metrics on compact Kahler manifolds*, Invent. Math. 81 (1985), **no. 2**, 359–372.
4. Chow, Bennet, *A Gradient Estimate for the Ricci Kähler Flow*, Anals of Global Analysis and Geometry 19 (2001) , 321–325.
5. Ecker, K. and Huisken, G., *Interior estimates for hypersurfaces moving by mean curvature*, Invent. Math. **105** (1991), 547–569.
6. Grigor'yan, A.A., *The heat equation on non-compact Riemannian manifolds*. Math. USSR Sbornik, Vol.**72** (1992), No.1.
7. Saloff-Coste, L., *Uniformly elliptic operators on Riemannian manifolds*, J. Differential Geom. **36** (1992), no. 2, 417–450.
8. Shi, Wan-Xiong, *Ricci Flow and the uniformization on complete non compact Kahler manifolds*, J. of Differential Geometry. **45** (1997), no. 1, 94–220.
9. Tian, G. and Yau S.T., *Existence of Kahler Einstein metrics on complete Kahler manifolds and their applications to algebraic geometry*, in "Mathematical Aspects of String Theory", edited by S.T. Yau, World Scientific, 1986.
10. Tian, G. and Yau S.T., *Complete Kahler manifolds with zero Ricci curvature. II.*, Invent. Math. **no. 106**(1990), no.1, 27–60.
11. Yau, S.-T., *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampre equation. I*, Comm. Pure Appl. Math. 31 (1978), no. 3, 339–411.
12. Yau, S.-T., *The role of partial differential equations in differential geometry*, Proceedings of the International Congress of Mathematicians (Helsinki, 1978), pp. 237–250, Acad. Sci. Fennica, Helsinki, 1980.
13. Yau, S.-T., *A general Schwarz lemma for Kähler manifolds*, Amer. J. Math. 100 (1978), no. 1, 197–203.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF BRITISH COLUMBIA,
ROOM 121, 1984 MATHEMATICS ROAD, VANCOUVER, B.C., CANADA V6T 1Z2
E-mail address: `chau@math.ubc.ca`

THE INSTITUTE OF MATHEMATICAL SCIENCES AND DEPARTMENT OF MATHEMATICS,
THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG,
CHINA.

E-mail address: `lftam@math.cuhk.edu.hk`